



## Toxicity and Ferti-irrigation Assessment of Paper Mill Effluent on Agronomic Characteristics of Black Gram

Akhil Gupta<sup>1</sup>, Aprajita Singh<sup>1</sup>, J. K. Pathak<sup>2</sup> & Anamika Tripathi\*

<sup>1</sup> Pollution Ecology Research Laboratory, Deptt. of Botany, Hindu college Moradabad, University of Rohilkhand- 244001, INDIA

<sup>2</sup>Hydrobiology lab, Deptt. of Zoology, Hindu college Moradabad, University of Rohilkhand.-244001, INDIA

\* Correspondance: E-mail: [akhil.gupta4@gmail.com](mailto:akhil.gupta4@gmail.com)

(Received 31 Jan, 2016; Accepted 05 Feb, 2016; Published 06 Feb, 2016)

**ABSTRACT:** Impact of paper mill effluent (PME) was analyzed on black gram (*Vigna mungo T-9*). Selected parameters selected were chlorophyll, protein content, root length, shoot length, leaf area and total biomass of *V. mungo T-9*. Effects of paper mill effluent were analyzed at different dilutions viz. 5%, 10%, 20%, 40%, 30%, 50%, & control. At more than 10% of effluent concentration decline was reported in biomass, protein and chlorophyll (4.8 mg/g highest). However, protein content was reported highest (23.6 mg/g) at 20% of concentration of effluent and showed decline further at higher concentration. Simultaneously soil samples were analyzed which showed, gradual increase in N, PO<sub>4</sub>, Na, K, Mg, and Ca at increasing concentration of paper mill effluent. Biochemical parameters like chlorophyll declined (2.4 mg/g) and protein declined (9 mg/g) clearly above twenty percent of effluent concentration which resulted in decrease in root length, shoot length, leaf area and biomass of *Vigna mungo T-9*.

**Keywords:** Effluent; Chlorophyll; Protein; Biomass ; leaf area; Soil and black gram.

**INTRODUCTION:** Pulp and paper in industry is the sixth largest polluter after oil, cement, leather, environment (Ali et al. 2001). Pulp and paper produced corresponds to only 40-45% of original weight of wood, the effluent are heavily loaded with organic matter the utility potential of paper mill effluent for irrigation of crop field has been a controversial proposition due to contradictory reports obtained on the effects of various effluents on crop plant response, (Narwal et al., 2005 and Madan, 2012). Effluents released from pulp and paper mill contains large amount of heavy metals which get accumulated in plant and soil. They cause huge damage to plants and biological systems (Chandra et al., 2010, Karthiwal, 2012 and Mehta and Bhardwaj, 2012) and even to ground water quality and soil (Balakrishnan et al., 2005, Senthilkumar et al., 2011, Chopra et. al., 2011, Tripathi et al., 2014) and even cancer risk to communities (Soskolne et al., 2010). In agriculture, irrigation water can affect soil characteristics and agricultural crop growth (Almodares and Shariff, 2007, Ebead et al., 2013). In some studies characteristics of effluents of industries and agronomic properties of various crops have been determined (Sawaf, 2005 and Mendoza et al., 2006, Bhargava et al., 2008, Iqbal et al., 2013). The black liquor from pulp and paper mill is a complex colloidal solution of various organic and inorganic polymeric substances like lignin, carbohydrate and their derivatives.

The volume and chemical nature of paper mill effluent depends on the type of manufacturing process adopted. Per kilogram of paper production discharges about 270-450 liters of effluent, along with approximate 50 g. of lignin. Contrary to this the small paper mills without soda recovery discharge 300-400 liters of black liquor effluent containing 200-250 gm. lignin per kg. (Garg and Modi, 1999). Biomethanated and partially treated effluent promotes growth in certain wetland plants (Singh et. al., 2012). *Vigna mungo T-9* commonly known as black gram is an important crop in Indian subcontinent. It is cultivated in tropical, semitropical temperate climate. It attains height between 25-50 cm and is highly rich in protein content. It is erected; bushy plant grows in warmer regions of India. The objective of the study is to analyze the effect of paper mill effluent on soil characteristics, morphological response, protein and carbohydrate content of *V. mungo var T-9* at different effluent concentrations.

**MATERIALS AND METHODS:** The paper mill effluent was collected near Shakumbhari Paper Mill Moradabad, situated at Moradabad-Agra highway. Immediately after collection, the effluent was brought to the laboratory with proper storage for analysis of physico-chemical parameters.

The physico-chemical parameters of paper mill effluent were analyzed by standard methods described by APHA (2005). Chemical properties of soil (taken in different pots) were analyzed by standard methods described by Trivedy and Goel (1986). Whereas heavy metals were analyzed by Inductively Coupled Plasma-Optical Emission Spectrophotometer after digestion process.

The pH of the sample was measured by calibrating digital pH meter (Systronics MK6) with buffer solution of pH 4 and 9. Turbidity was measured by turbidity meter using the technique of nephelometry, which measures the amount of light scattered at right angles to an incident light beam by particles present in a fluid sample (Sadar 1996).

Total dissolved solids (TDS) of the sample were determined by Hanna Pocket TDS meter. (HI -98130). Conductivity of the effluent sample was measured with a probe and conductivity meter. A voltage is applied between the two electrodes in the probe immersed in the sample water. The drop in voltage caused by the resistance of the water is used to calculate the conductivity per centimeter, as per the manual, by using potassium chloride and distilled water. Dissolved Oxygen is the oxygen concentration in the sample which is defined as the number of milliliters of dioxygen gas (O<sub>2</sub>) per liter of water. DO in effluent sample was measured by Winklers titration. Biological Oxygen Demand (BOD) was measured after five days BOD at 20°C was determined by standard dilution technique according to standard methods. Chemical Oxygen Demand (COD) was measured by Potassium dichromate reflux method.

Nitrate Determination is done by spectrophotometer at a wavelength of 480 nm Cataldo et al., (1975). Inorganic Phosphate estimation in effluent and soil was done by Vanado Molybdophosphoric Acid Method. Ammonia of the effluent was determined as Total Kjeldal Nitrogen (TKN) by Kjeldal method (1883). Chlorides of the samples were estimated by silver nitrate solution, Sodium, Potassium, Calcium and Magnesium in effluent and soil were measured by Flame Photometer. Alkalinity of the sample was measured by EDTA titration method. Salinity of the effluent was measured by salinity meter Wattson (WS-SAs 287). Total hardness was estimated by EDTA titration. For heavy metal i.e. Cu, Cd, Cr, Pb and Zn in effluent and soil samples were analysed by ICP-OES spectrophotometer.

In order to study the effect of paper mill effluent on *V.mungo* growth 21 pots of 20×16 cm height were filled with approximate 2 kg of garden loam soil. Seeds were sterilized by 0.1% mercuric chloride and

soaked and germinated. Ten samplings were sown in each pot. The pots were irrigated with different concentration (0.00%, 10%, 20%, 30%, 40%, and 50%) of effluent. These pots were irrigated by 250 ml of effluent of each concentration with 2 day intervals throughout the study period. Seeds sown in pots filled by garden soil and irrigated with sterilized tap water served as control. Regarding germination of seeds emergence of radical as noted as criterion of germination of seed. Number of seeds germinated during stipulated time frame was expressed as percent of germination. After 30 days 3 plants were taken from each set and washed twice by sterilized water. Shoot and root length were measured by meter scale. Biomass of plant was determined by dry weight method as described by APHAs (2005), where plant after harvesting, plant roots, shoots and leaves were separated and dried at 65 C° in hot air oven for 48 hours. The difference of fresh weight and dry weight was recorded as biomass of plants.

#### Analysis of morphological and biochemical parameters:

For pigment determination, 500 mg of dry leaf were homogenized in 20 ml of 80% acetone using mortar and paste and centrifuged at 6000×g for 15 minutes finally the supernatant was made up to 20 ml and Optical Densities (O.D.) were measured at 480 and 510 nm wavelength for carotenoides and 645 nm and 663 nm for chlorophyll on a UV-VIS spectrophotometer (Systronics Model 119, India). The amount of chlorophyll a and b and carotenoid were calculated by using the formulae give by Machlachan and Zalick 1963 and Duxbury and Yentsch (1956) respectively.

$$\text{Chlorophyll a} \left( \frac{\text{mg}}{\text{g}} \text{ dry leaf} \right) = \frac{[12.3 \times D_{663} - 0.86 \times D_{645}] \times V}{d \times 1000 \times w}$$

$$\text{Chlorophyll b} \left( \frac{\text{mg}}{\text{g}} \text{ dry leaf} \right) = \frac{[19.3 \times D_{645} - 3.6 \times D_{663}] \times V}{d \times 1000 \times w}$$

$$\text{Total Chlorophyll} \left( \frac{\text{mg}}{\text{g}} \text{ dry leaf} \right) = \text{Chlorophyll a} + \text{Chlorophyll b}$$

Where; V = Volume of extract (ml), d= length of light path (cm), w = dry weight of leaf.

Protein content was estimated by Lowry et al. (1951). Testing was done as per the manual and absorbance was measured at 660 nm against blank. Then standard curve was made by taking concentration of protein along X-axis and absorbance at 660 nm along Y-axis.

While Leaf area of black gram seedlings was measured by leaf area meter (Systronics 128).

**RESULTS AND DISCUSSION:** It is evident from Table 3, that paper mill effluent was alkaline in nature having pH 7.4. The values of TDS, BOD, COD exceeded the permissible limits of ISI 1982 (IS: 2296), Observed BOD were very high, 295 mg/l (Table 1). As compared to the normal values this indicates the high concentration of organic material in effluent. This work is also supported by the work of (Ningomban et al., 2009, Medhi et al., 2011 and Madan et al., 2012). COD is more realistic parameter which indicates the pollution status of water body. COD values were quite high 700 mg/l (Table 1) As per Indian standards maximum values of COD for pulp and paper mill effluent is 300mg/l. Phosphate, Nitrate and Total Kjeldhal Nitrogen were also found high (710, 357, 487 mg/l respectively).

Heavy metals in the effluent were found to be exceeding the permissible limits (Table 1) Cr, Cu, Zn, and Cd and Fe were found to be 0.59 mg/l, 0.64 mg/l, 1.43 mg/l, and 0.10 mg/l respectively. Since the effluent is rich in heavy metals, the growth of *Vigna* is affected, which are irrigated by this effluent at different concentration (Table 5). These results of the physico-chemical analysis coincide with earlier reports (Madan 2012). Effluent loaded with heavy metals leads to increase in concentration of metals in soil and vegetation (Satpati et al., 2010). Moreover, heavy metals accumulation have adverse impact on chlorophyll content on various vegetables, crops and alters the biomass production and yield (Sharma et al., 2008, Medhi et al., 2011). But, at the same time, certain plants which accumulate heavy metals in greater quantities can serve as source of phytoremediation (Rai, 2010).

Electrical conductivity is highly correlated with phosphate ( $r = 0.967$ ), Mg ( $r = 0.953$ ), Ca ( $r = 0.904$ ), while carbon is highly correlated with organic matter ( $r = 0.876$ ) present in effluent. Mg & EC ( $r = 0.953$ ) were found to have high correlation value. Mg & Ca ( $r = 0.906$ ) were highly correlated, while Na & Mg ( $r = 0.454$ ) were found to show lesser degree of correlation. Carbon was found to be negatively correlated with Nitrogen ( $r = -0.88$ ). Nitrogen and Potassium showed lesser degree ( $r = 0.044$ ) degree of correlation. Golla *et al.*, (2012) reported similar findings with cotton ginning mill effluent.

From Table 2, it is evident that pH of soil increased with increasing concentration due to presence of high amount of  $CO_3$  and Ca in paper mill effluent. Madan, (2012). In present study, organic matter, Ca, Mg, K, Na concentration decreased with increasing the efflu-

ent concentration. Organic matter was found high in soils irrigated with 10% of effluent concentration. Sodium values were found highest in soils irrigated with 50 percent of effluent concentration. Tripathi et al., (2014) has revealed that soils irrigated with paper mill effluent have high microbial diversity both in term of structure and function than the control soil.

**Table 1: Mean value of different parameters (mg/l) after effluent which was used in different dilutions. Mean value along with Standard Deviation, n=5.**

Parameters	Average	SD
pH	7.42	0.371
EC (mmho/cm)	1506.2	58.486
TDS	1311.4	30.342
DO	1.26	0.242
BOD	295.4	19.946
COD	700.4	24.303
Na	304.2	39.555
K	278	67.814
Chloride	647.8	87.303
Alkalinity	702.6	28.204
Nitrate	357.4	24.532
TKN	487.6	151.550
Salinity	0.68	0.147
Turbidity	66.8	2.135
Phosphate	710.6	100.039
Cu	0.64	0.196
Zn	1.5	0.963
Pb	0.56	0
Hg	0.00	0.000
Cd	0.108	0.048
Cr	0.59	0.341

At more than 10% effluent concentration, test plant showed inhibitory effect on growth on roots shoots length biomass. Similar type of tolerance studies were carried out on seven varieties of finger millet to factory effluent. The varietal screening of groundnut for tolerance to fertilizer factory effluent (Sundaramoorthy, 1995) and sugar mill effluent (Ezhumalai 2002 and Sundai, 2003) were reported. Similarly, the effect on paddy cultivars for tolerance to tannery effluent and Sugar mill effluent (Sundaramoorthy et al., 2003) was reported High amount of chemicals and salt and high concentration of sodium increases the exchangeable sodium percentage of the soil to high level of phytotoxicity. The maximum growth of *V. mungo* was noted upto 10% of effluent. Coefficient of correlation analysis and regression analysis is given in figure one and two.

Compared to control, total chlorophyll content was found to be most sensitive parameter. Further the metals present in paper mill effluent generate active oxygen species and reduced total chlorophyll may be due to phytotoxic consequences of reactive oxygen species (Chandra et al., 2010). Reduced protein content in above 20% effluent irrigated plants may also be attributed to the presence of toxic concentration of heavy metals in effluent. However, the increase in Protein content in *V. mungo* exposed to low concentration of paper mill effluent may be due to synthesis of stress protein. Bharagava et al., (2008) reported that distillery effluent irrigation increase chlorophyll and protein contents in Indian mustard plants (*Brassica nigra* L.) at 25 and 50% sugar mill effluent concentrations followed by decrease in 75% and 100% sugar mill effluent.

The study revealed that at more than 10% concentration of effluent all the growth parameters showed

decline except protein content. The improvement of vegetative growth may be attributed to the role of potassium in nutrient and sugar translocation in plants and turgor pressure in plant cells. It is also involved in cell enlargement and in triggering young tissue or meristematic growth. (Kumar, 2012) Chlorophyll content was higher due to use of 20% dilution of paper mill effluent. This contains optimum contents of nutrients required for maximum vegetative growth of *Vigna mungo T-9*. This coincides with the results of Medhi et al., (2011), Gotosa et al., (2011), Madan et al., (2012) and Kumar et al., (2013), Mishra et al., (2013) regarding paper mill effluent. Along with this analysis of soil samples irrigated by different effluent concentration have shown that effluent increases the soil mineral content and organic matter in small quantities. From the above results it can be concluded that, effluent can be used safely for *V. mungo var T-9* cultivation only after proper dilution up to certain extent

**Table 2: Values of different soil parameters (mg/l)s at different concentrations of effluent.**

PME Conc. Per 2kg	pH	EC (mmho/cm)	C%	Organic Matter	N%	PO <sub>4</sub> mg/l	Na	K	Mg	Ca
Before PME Application	6.2	0.345	0.42	0.81	0.08	0.8	21.7	3.6	11.7	75
10%	6.3	0.346	0.59	1.07	0.03	0.81	16.6	1.5	13.7	76
20%	6.5	0.355	0.52	0.89	0.03	0.97	17.5	2.2	14.1	77
30%	6.9	0.359	0.53	0.91	0.03	1.31	20.5	3.7	15.2	88
40%	7.3	0.378	0.57	0.93	0.02	1.77	22.5	4.7	16.4	95
50%	7.4	0.396	0.59	0.96	0.03	1.91	22.6	5.3	18.5	95

**Table 3: Correlation among different parameters in soil analysis.**

	Ph	EC	C%	OM	N%	PO <sub>4</sub>	Na	K	Mg	Ca
pH	1									
EC	0.947	1								
C%	0.583	0.56	1							
OM	0.155	0.152	0.876	1						
N%	-0.608	-0.488	-0.886	-0.665	1					
PO <sub>4</sub>	0.993	0.967	0.547	0.124	-0.53	1				
Na	0.652	0.647	-0.142	-0.455	0.18	0.707	1			
K	0.819	0.824	0.081	-0.298	-0.044	0.86	0.961	1		
Mg	0.947	0.953	0.743	0.367	-0.694	0.938	0.454	0.675	1	
Ca	0.988	0.909	0.544	0.139	-0.558	0.983	0.706	0.845	0.906	1


**Table 4: Values of chlorophyll and protein content (mg/gm) of *V. mungo* at different concentrations of Paper Mill Effluent.**

Effluent	Chl. mg/g	Protein
0%	4.4±.21	18.46
5%	4.7±.08	18.5±.5
10%	4.8±.16	19.7±.5
20%	3.6±.205	23.6±1.5
30%	3.1±.04	18.7±1.5
40%	2.6±.12	12.2±1.5
50%	2.4±0	9.0±1.2


**Table 5: Root length, shoot length and biomass of *V. mungo* at different concentrations of effluent.**

Effluent Concentration (%)	Shoot Length (cm)	Root Length (cm)	Biomass Per Plant (gm)
0.0	24.2±.25	5.6±.05	0.264±.005
5%	24.7±.25	5.5±.4	0.283±.003
10%	25.6±.4	6.00±.1	0.317±.007
20%	25.1±.85	6.4±.05	0.258±.0315
30%	22.2±1.2	5.9±.4	0.23±.003
40%	21.5±.5	5.8±.1	.225±.025
50%	20±1	5.6±.1	0.17±.025

Thus it can be concluded that at lower concentration effluent increases the plant growth (Rana et al., 2012 and Iqbal et al., 2013.) At more than 20% effluent concentration, test plant showed inhibitory effect on growth on roots shoots length biomass. Further the metals present in paper mill effluent generate active oxygen species and reduced total chlorophyll may be due to phytotoxic consequences of reactive oxygen species (Chandra et al., 2010).



**Figure 1: Coefficient of correlation between protein content of *Vigna mungo* var *T-9* and different dilutions of effluent.**



**Figure 2: Coefficient of correlation between chlorophyll content of *Vigna mungo* var *T-9* at different dilutions of effluent.**

Reduced protein content in above 20% effluent irrigated plants may also be attributed to the presence of toxic concentration of heavy metals in effluent. However, the increase in Protein content in *V. mungo* exposed to low concentration of paper mill effluent may be due to synthesis of stress protein. Biochemical parameters, chlorophyll and protein declined clearly above twenty percent of effluent concentration which resulted in decrease in root length, shoot length leaf area and biomass of *Vigna mungo T-9*.

**CONCLUSION:** Since industrial effluent poses a ecological risk. Phytotoxic effects of paper mill effluent and ferti-irrigation studies of effluent with black gram and chlorophyll and protein analysis of it in black gram are highly valuable since *Vigna mungo T-9* is prominent pulses of north India. Further, incorporation with one other remedial technique such as soil amendments, sprinkler irrigation, drip irrigation and intercropping system can reduce the phytotoxic effects of effluent and can improve ferti-irrigation potential of industrial effluent as shown by *Vigna mungo T-9* at optimum level of dilution.

**ACKNOWLEDGEMENT:** One of the authors, Akhil Gupta (SRF, NET) expresses his sincere thanks to UGC for providing the necessary financial assistance for conducting the work. Authors are highly thankful to Research testing and calibration laboratory. Ministry of textiles, Govt. of India Moradabad for providing us cooperation during testing and research work. Authors are also highly thankful to Dr. V. K. Vats, Head Department of Botany, Hindu college, Moradabad, for his invaluable guidance during the study.

## REFERENCES:

1. APHA., (2005). Standard methods the examination of water and wastewater, 21<sup>st</sup> Edn., APHA, AWWA WPCF, Washington DC USA.
2. Almodares A. & Sharif M. E. (2007) Effects of irrigation water qualities on biomass and sugar contents of sugar beet and sweet sorghum cultivars, *Journal of Environmental Biology*, 28, 213-216.
3. Ali M. and Sreekrishnan T. R. (2001) Aquatic toxicity from pulp and paper mill effluents; a review, *Advances in Environmental Research*, Elsevier Publications, 5, 175-196
4. Balakrishnan V., and Karruppusamy S. (2005) Physico chemical characteristics of drinking water samples of pilani, Tamilnadu, *Journal of Ecotoxicology and Environmental Monitoring*, 15, 223-35.
5. Bharagava R. N., Chandra R., and Rai V. (2008) Phytoextraction of trace elements and physiological changes in Indian mustard plants (*Brassica nigra* L.) grown in posts methanated distillery effluent (PMDE) irrigated soil, *Bioresource Technology*, 99, 8316–8324.
6. Chandra R., Bhargava R.N., Yadav S., and Mohan B. (2009) Accumulation of heavy metals in wheat (*Triticumaestivum*) and Indian mustard (*Bassicajuncea*) irrigated with distillery and tannery effluents, *Journal of Hazardous Matter.*, 169, 1514-1521.
7. Chandra R. P., Abdulsalam A. K., Salim N. A. and Puthur J. T. (2010) Distribution of bio accumulated cadmium and chromium in two *Vigna species* and associated histological variations, *Journal of Stress Physiology and Biochemistry*, 6, 4-14.
8. Chopra A. K., Srivastava K. V. (2011) Study of agro potentiality of paper –mill effluent and synthetic nutrient (DAP) on *Vigna unguiculata* L.(walp) cowpea, *Journal of Chemical and Pharmaceutical Research*, 3, 151-165.
9. El-Sawaf, Nadia. (2005) Response of Sorghum spp. To sewage wastewater irrigation, *International Journal of Agriculture Biology*, 7, 869–874.
10. Ezhumalai D. (2002) Screening of black gam variety of tolerant to sugar mill effluent treatment Mphil thesis to annamalai university
11. Garg S. K. and Modi D. R. (1999) Decolorisation of pulp and paper mill effluent by white rot fungi. *Cit,Rev, Biotechnol*, 19, 85-112.
12. Gotosa J., Nezandonyi K., Kanda A., Mushiri S. M., Kundhlande A., and Nyamunungure T. (2011) Effects of irrigating Eucalyptus grandis plantation with a mixture of domestic and pulp and paper mill effluent on soil quality at a site in northern Zimbabwe, *J. of sustainable development in Africa*, 13, 152-55.
13. Gharoobi B., Ghobani M., and Nehzad M. G. (2012) Effects of different levels of osmotic potential on germination percentage and germination ate of barley, corn and canole, *Iranian j. of Plant Physiol.*, 2, 413-417.
14. Iqbal S., Younas U., Chan K. W., Saeed Z., Shaheen M. A., Akhtar N., and Majeed A. (2013) Growth and antioxidant response of *Brassica rapa* var. *rapa* L. (turnip) irrigated with different compositions of paper and board mill (PBM) effluent, *Chemosphere*, 91, 1196-1202.
15. ICMR (1974) Tolerance limits for industrial effluents discharged on land for irrigation, IS-3307 New Delhi.
16. Kathirvel P. (2012) The effect of dye factory effluent on growth, yield and biochemical attributes of Bengal gram (*Cicer arietinum*), *Int. J. App. Biol. Pharmace. Technol.*, 3, 146-150.
17. Kumar V., Chopra A. K., and Pathak C. (2010) Agropotentiality of paper mill effluent on characteristics of *Trigonella foenum-graccum*, Dept of zoology and environmental sciences. Gurukul Kangri University Haridwar, *Newyork Science Journal*, 3, 57-63.
18. Kumar V, and Chopra A.K. (2013).Ferti-irrigation effect of paper mill effluent on agronomical characteristics of *Abel moschus esculentus*(okra), *Pakistan J. of Biological science*,16, 1426-1437.
19. Kumar V., Dhall P., Kumar R., Singh Y. P., Kumar A. (2012) Bioremediation of agro-based pulp and paper mill effluent by microbial consortium comprising Antochthonous bacteria, *The Scientific World Journal*, Article ID 127014,doi;10;1100/2012/12701
20. Lngle, S. T. (2000) Pollution potential of pulp and paper mill effluent, *Bull. Environ.Sci.*, 18, 21-24
21. Madan S., and Pertibha (2013) Morphological response of *Cicerarietinum* exposed to paper mill effluent, *J. of Environ. Biol.*, 34, 779-782.
22. Malla L, and Mohanty B. K., (2005) Effect of paper mill effluent on germination of green gram and growth behavior of its seedling, *J. of Environ.Biol*, 26, 379-382.
23. Mehta A., and Bhardwaj N. (2012) Phytotoxic effects of industrial effluents on seed germinationand seedling growth of *Vigna radiata* and *Cicer arietinum*. *Global J.Biosci, Biotechnol*, 1-5.
24. Mendoza J., G. Tatiana, C. Gabriela, and S. M. Nilsa. (2006) Metal availability and uptake by sorghum plants grown in soil amended with sludge from different treatments, *Chemosphere*, 65, 2304–23012.

25. Medhi U. J, Talukdar A. K., and Deka S. (2011) Impact of paper effluent on growth and development of certain agricultural crops, *Environmental Biology*, 32, 185–188.
26. Mishra, S., Mohanty M., Pradhan, H., Patra K., Das R. and Sahoo S., (2013) Physico-chemical assessment of paper mill effluent and its heavy metal remediation using aquatic macrophytes—a case study at JK Paper mill, Rayagada, *India Environ Monit Assess* 185:4347–4359 DOI 10.1007/s10661-012-2873-9
27. Narwal R. P., Singh A. and Dahiya S. S., (2005) Effect of paper mill effluent's irrigation on soil and plants. Int. conf. on Energy, Environment and Disaster. INCEED 2005, Charlotte NC USA.
28. Ningomban L. D., Yadav I. C., Shihua I. I., Singh S., and Belagali S. L., (2011) Physiochemical characteristics of paper industry effluents - a case study of south India paper mills, *Environ Monit Assess*, 177;23-33. DOI 10.1007/10661-010-1614-1.
29. Rai K. P., (2010) Phytoremediation of heavy metals in a tropical impoundment of industrial region, *Environmental Monitoring and Assessment*, 148, 75-84.
30. Rana I, Dulaini A. L., Ismail N. B., Ibrahim M. H. (2012) The effect of industrial waste water in seed growth rate; a review, *International Journal of Scientific and Research Publication*, 2, 28-35.
31. Satpati S., Nayak S., and Gorai D. (2010) Effects of waste water irrigation on vegetables in relation to bioaccumulation of heavy metals and biochemical changes, *Environmental monitoring Assessment*, 16, 169-177
32. Sawaf E, Nadia (2005) Response of Sorghum spp. to sewage wastewater irrigation, *International Journal of Agriculture Biology*, 7, 869–874.
33. Senthilkumar D., Satheeshkumar P., and Gopalakishnan P. (2011) Ground water quality and assessment of paper mill effluent irrigated area using multivariate statistical analysis, *World applied sciences journal*, 13, 829-836.
34. Sharma B. and Chettri, M. K. (2008) Impacts of heavy metals on accumulation of essential micronutrients and chlorophyll in some vegetables, *Pollution research*, 27, 355-363.
35. Sharma R. K., Agrawal M. and Marshall F. M., (2004) Effects of waste water irrigation on heavy metal accumulation in soil and plants”, Paper presented in national seminar, Bangalore University Bangalore Abst. no. 7, 8.
36. Singh P. K., K. P. Sharma. (2012) Biomethanated distillery waste utilization by wetland plant; A new approach, *World Journal of Agricultural Science*, 8, 96-103.
37. Soskolne, C. L., and Sieswada E. L. (2010) Cancer risk associated with pulp and paper mills; A review of occupational and community epidemiology, *Chronic dis. Can.*, 29, 89-100
38. Sundaramoorthys P., J. Kunthijapatham (2001) Effect of paper mill effluent on seed germination and seedling growth of six varieties of ground nut, *Ecotoxicol Monitoring*, 10, 53-57.
39. Talukdar A. K, Medhi U. J., and Deka S. (2011) Impact of paper mill effluents on growth and development of certain agricultural crops, *J. of Environ Biology*, 32, 185-188.
40. Tamot S. & Sharma P. (2006) Physico-chemical status of upper lake (Bhopal India), Water quality with special references to phosphate and nitrate concentration and their impact on lake ecosystem, *Asian J. Exp. Sci.*, 20, 151-158.
41. Tripathi B. M., Kumari P, Weber K. P., Saxena A. K., Arora D. K., and Kaushik R. (2014) Influence of long term irrigation with pulp and paper mill effluent on the bacterial community structure and catabolic function of soil, *Indian J. Microbiol.*, 54, 65-73.